

INVENTORYING THE STELLAR INITIAL MASS FUNCTION OF EARLY-TYPE GALAXIES

C. TORTORA¹, A.J. ROMANOWSKY², AND N.R. NAPOLITANO³

¹Universität Zürich, Institut für Theoretische Physik, Winterthurerstrasse 190, CH-8057, Zürich, Switzerland

²University of California Observatories, 1156 High Street, Santa Cruz, CA 95064, USA

³INAF – Osservatorio Astronomico di Capodimonte, Salita Moiariello, 16, 80131 - Napoli, Italy

submitted to The Astrophysical Journal Letters, 19 July 2012

ABSTRACT

Given a flurry of recent claims for systematic variations in the stellar initial mass function (IMF), we carry out a novel inventory of the observational evidence using different approaches. This includes literature results, as well as our own new findings from combined stellar-populations synthesis (SPS) and Jeans dynamical analyses of data on ~ 4500 early-type galaxies from the SPIDER project. We focus on the mass-to-light ratio mismatch relative to the Milky Way IMF, δ_{IMF} , correlated against the central stellar velocity dispersion, σ_* . For the SPIDER sample, we find a strong correlation between δ_{IMF} and σ_* , independently of assumptions on the dark matter (DM) profile. The overall normalization of δ_{IMF} , and the detailed DM profile, are less certain, but the data are consistent with standard cold-DM halos, and a central DM fraction that is roughly constant with σ_* . For a variety of related studies in the literature, using SPS, dynamics, and gravitational lensing, similar results are found. Studies based solely on spectroscopic line diagnostics agree on a Salpeter-like IMF at high σ_* , but differ at low σ_* . Overall, we find that multiple independent lines of evidence appear to be converging on a systematic variation in the IMF, such that high- σ_* early-type galaxies have an excess of low-mass stars relative to spirals and low- σ_* early-types. Robust verification of super-Salpeter IMFs in the highest- σ_* galaxies will require additional scrutiny of scatter and systematic uncertainties. The combined implications for the distribution of DM are still inconclusive.

Subject headings: galaxies: evolution — galaxies: general — galaxies: elliptical and lenticular, cD

1. INTRODUCTION

The stellar Initial Mass Function (IMF) is fundamentally important to understanding both stellar populations and galaxies. The Milky Way (MW) IMF was originally characterized as a power-law mass-distribution, $dN/dM \propto M^{-\alpha}$, with $\alpha \sim 2.35$ (Salpeter 1955), and subsequently refined to flatten at lower masses ($M \lesssim 0.5M_\odot$; Kroupa 2001; Chabrier 2003).

Whether or not the MW IMF describes stellar populations elsewhere in the Universe cannot yet be said through direct star counts. There have been some indirect observational hints of IMF variations, and ample theoretical motivation for these, but no broadly convincing evidence has emerged (cf. Bastian et al. 2010).

This situation has recently changed, with a flurry of studies of early-type galaxies (ETGs; ellipticals and lenticulars) turning up indirect evidence for systematic IMF variations. These studies use models of stellar population synthesis (SPS) to fit integrated-light data (broad-band colors and spectroscopic features), and fall into two broad categories: “pure” SPS, and “hybrid” SPS+gravitating mass analyses.

The pure analyses rely on spectral lines that are differentially sensitive to giant or dwarf stars. These include the TiO feature at 6130 Å, the Na I doublet near 8190 Å, the Ca II triplet near 8600 Å, the Wing–Ford [Fe/H] band at 9915 Å, and the Ca I line at 10345 Å (e.g., Cenarro et al. 2003; van Dokkum & Conroy 2010; Spiniello et al. 2012; Conroy & van Dokkum 2012, hereafter CvD12; Ferreras et al. 2012; Smith et al. 2012).

The hybrid analyses assume an IMF and infer a stellar mass-to-light ratio Υ_* using a more conventional SPS

approach based on colors, and age- and metallicity-sensitive spectral lines. Estimates of Υ_* are also derived using dynamics or gravitational lensing. Comparison of the independent results then yields a revised IMF (e.g., Renzini & Ciotti 1993; Cappellari et al. 2006, 2012; Ferreras et al. 2008; Tortora et al. 2009, 2010; Grillo & Gobat 2010; Treu et al. 2010; Napolitano et al. 2010, 2011; Auger et al. 2010; Grillo 2010; Thomas et al. 2011; Spiniello et al. 2011; Dutton et al. 2012b,a; Sonnenfeld et al. 2012; Wegner et al. 2012).

One may characterize a revised IMF through its Υ_* relative to a MW-disk IMF, $\delta_{\text{IMF}} \equiv \Upsilon_*/\Upsilon_{*,\text{MW}}$ (the “mismatch parameter”), where for reference we adopt the Chabrier IMF. Remarkably, almost all the above studies found “heavy” IMFs ($\delta_{\text{IMF}} \gg 1$) for the most massive ETGs. Less massive ETGs, and spiral galaxies, appear to have “normal/light” IMFs ($\delta_{\text{IMF}} \lesssim 1$; e.g., Bell & de Jong 2001; Bershadsky et al. 2011; Suyu et al. 2012; Brewer et al. 2012).

These IMF findings are both potentially revolutionary, and highly controversial, and demand further investigation. In particular, with hybrid analyses there are lingering questions about degeneracies associated with the distribution of nonbaryonic dark matter (DM). The time is also ripe to inventory the results to date, and see if the apparent emerging consensus holds up under quantitative, systematic comparison – which could provide pressing motivation for understanding the physical origins of the trends. Comparisons were made for some hybrid analyses (Thomas et al. 2011; Dutton et al. 2012a; Wegner et al. 2012), but not for the pure SPS work.

In this Letter we carry out such an inventory, while presenting our own novel results for a large sample of ETGs

for reference, following the dynamical+SPS analyses of Tortora et al. (2012, hereafter T+12). We focus on the trends in δ_{IMF} with central stellar velocity dispersion, σ_* . σ_* is widely considered as crucially connected to galaxy evolution, and unlike Υ_* , is relatively independent of the bandpass, and of δ_{IMF} .

We describe our data and analysis methods in Section 2. We present our results in Section 3 and make literature comparisons in Section 4. We summarize the conclusions and outlook in Section 5.

2. DATA AND ANALYSIS METHODS

We apply a combination of SPS and stellar dynamical models to a sample of ~ 4500 giant ETGs, in the redshift range of $z = 0.05\text{--}0.1$, drawn from the Spheroid’s Panchromatic Investigation in Different Environmental Regimes project (SPIDER; La Barbera et al. 2010). The details of sample selection and methods can be found in T+12, and here we provide only an overview of our two Υ_* -estimation methods.

Our data include optical+near-infrared photometry [from the Sloan Digital Sky Survey (SDSS) and the UKIRT Infrared Deep Sky Survey-Large Area Survey]¹, high-quality measurements of galactic structural parameters (effective radius R_e and Sérsic index n), and SDSS central-aperture velocity dispersions σ_{Ap} . The SPS-based Υ_* values were derived by fitting Bruzual & Charlot (2003) models to the multi-band photometry, assuming a Chabrier IMF.

Our dynamical-mass estimates use spherical isotropic Jeans equations fitted to the σ_{Ap} data. We also extend the T+12 analysis using two-component mass models: a Sérsic-based stellar distribution following the K -band light, and a standard DM profile. The data do not constrain both components simultaneously, so instead we adopt a series of plausible assumptions about the DM (cf. Cappellari et al. 2012).

Our DM models are based on the Navarro et al. (1996, hereafter NFW) profile, with an adjustable degree of baryon-induced adiabatic contraction (AC). For the virial mass and concentration (M_{vir} , c_{vir}), we adopt mean trends for a WMAP5 cosmology (Macciò et al. 2008). We take an M_{vir} –stellar-mass (M_*) relation from Moster et al. (2010, M+10 hereafter). Each galaxy’s mass model then has one free parameter, Υ_* , plus optional AC (Gnedin et al. 2004, G+04 hereafter), providing our no-AC-NFW and AC-NFW base-models.

We explore the sensitivity of our results to these assumptions by doing the analyses with the following alternatives: a) AC recipes of varying strengths (Blumenthal et al. 1986; Abadi et al. 2010); b) M_{vir} with a fixed value for the entire sample: $M_{\text{vir}} = 10^{12}M_\odot$, $10^{13}M_\odot$, or $10^{14}M_\odot$; c) WMAP3-based $c_{\text{vir}}\text{--}M_{\text{vir}}$ relation (Macciò et al. 2008); d) no DM is present; e) mild kinematic anisotropy, with $\beta = -0.2$ or $+0.2$; f) $c_{\text{vir}}\text{--}M_{\text{vir}}$ relation altered to mimic a warm dark matter (WDM) cosmology, assuming different particle masses (Schneider et al. 2012, S+12 hereafter).

In order to study the mean trends of Υ_* with velocity dispersion, we construct “average” galaxies from the data. Dividing our sample into 10 σ_{Ap} -bins, for which we compute median values of M_* , R_e , n , R_{Ap} , and σ_{Ap} .




Figure 1. Galaxy sample properties, binned by velocity dispersion: physical aperture radius, SPS-based stellar mass (Chabrier IMF), effective radius, and Sérsic index. The points and error-bars show the medians and 25–75 percentile scatter.

We show these values, and their 25–75 percentile scatter, in Fig. 1.

Our final analysis-products, for each galaxy bin and mass model, will be the SPS-determined $\Upsilon_{*,\text{MW}}$, the dynamically-determined Υ_* , the inferred δ_{IMF} , and the inferred central DM-fraction, $f_{\text{DM}} \equiv 1 - \Upsilon_*/\Upsilon_{\text{dyn}}$. For homogeneity, we convert σ_{Ap} to σ_e (the value at R_e), using the best-fitting relation in Cappellari et al. (2006) (also done for the literature results later).

3. RESULTS: IMF AND DM FRACTION TRENDS

Our main IMF results from SPIDER are shown in Figure 2. The two thick black curves correspond to our standard no-AC-NFW (solid) and AC-NFW (long-dashed) models, and the suite of alternative models are also plotted, as labeled.

It is clear that the overall normalization of δ_{IMF} is degenerate with the adopted halo model, as DM can be traded against stellar mass. However, for a given flavor of halo model, there is always a strong correlation between δ_{IMF} and σ_* (the $\delta_{\text{IMF}}\text{--}M_*$ trend is weaker). We quantify this correlation for each model with a log-log fit, reporting the best-fit parameters in Table 1; the typical relation is $\delta_{\text{IMF}} \propto \sigma_e$.

Other models of potential interest are low-density DM cores (Burkert 1995), and alternatives to DM (e.g., Milgrom 1983). These would imply similar $\delta_{\text{IMF}}\text{--}\sigma_e$ slopes to our limiting no-DM model.

One could alter these slopes with additional model-tuning; e.g., with strong-AC at high- σ_e and halo *ex*-

¹ <http://www.sdss.org>, <http://www.ukidss.org>




Figure 2. IMF mismatch-parameter, $\delta_{\text{IMF}} = \Upsilon_*/\Upsilon_{*,\text{MW}}$, vs. velocity dispersion σ_e , for the SPIDER sample. *Top*: results for several different mass models. Horizontal lines show reference values for Salpeter, Kroupa, Chabrier IMFs (top-to-bottom). *Bottom*: residuals relative to the fiducial no-AC-NFW result, for an expanded series of mass models (see legend at right, explanation in the main text, and reference list for abbreviations). The overall normalization of δ_{IMF} depends on the model assumptions, but a steep relation between δ_{IMF} and σ_e emerges as a robust result.

Table 1

Best-fit parameters for the relation $\log \delta_{\text{IMF}} = a + b \log \frac{\sigma_e}{200 \text{ km/s}}$, for model suite (see main text).

Model	a	b
$M_{\text{vir}} - M_*$ (M+10)	0.18	0.86
$M_{\text{vir}} - M_*$ (M+10) + AC (G+04)	0.11	1.07
$M_{\text{vir}} - M_*$ (M+10) + AC (A+10)	0.16	0.93
$M_{\text{vir}} - M_*$ (M+10) + AC (B+86)	0.04	1.29
$M_{\text{vir}} = 10^{12} M_{\odot}$	0.19	0.88
$M_{\text{vir}} = 10^{13} M_{\odot}$	0.18	0.97
$M_{\text{vir}} = 10^{14} M_{\odot}$	0.16	1.09
$M_{\text{vir}} - M_*$ (M+10) - WMAP3	0.18	0.85
no DM	0.21	0.79
$\beta = +0.2$	0.17	0.87
$\beta = -0.2$	0.19	0.88
CDM (S+11)	0.17	0.90
$m_{\text{WDM}} = 1 \text{ keV}$ (S+11)	0.18	0.88
$m_{\text{WDM}} = 0.5 \text{ keV}$ (S+11)	0.18	0.84
$m_{\text{WDM}} = 0.25 \text{ keV}$ (S+11)	0.19	0.80

pansion at low- σ_e , one could completely flatten out the trend. We currently have no a priori motivation for such variations, but they would be a way to preserve IMF universality.

We next examining the overall IMF normalization, with the no-AC and AC-NFW cases bracketing the most plausible range of models. For reference, we show δ_{IMF}

predictions for several standard IMFs (Salpeter, Kroupa, Chabrier). We note that at a fixed IMF, age, and metallicity, the Υ_* and δ_{IMF} values are uniquely predicted, but the reverse is not true. A given δ_{IMF} result can imply multiple IMF solutions, particularly if one allows for mass-functions more complicated than a pure power-law.

If we adopt a no-AC-NFW model, a MW-like IMF is implied for low- σ_e galaxies, and a Salpeter IMF for high- σ_e . For AC-NFW, the low- σ_e galaxies have sub-MW IMFs, and the high- σ_e ones have IMFs intermediate to Kroupa and Salpeter.

In all cases, extremely bottom-heavy IMFs ($\alpha \gtrsim 2.6$) are ruled out, on average. Even if one assumed *no* DM, such IMFs would violate the dynamical constraints on the overall mass-to-light ratio.

Although resolving the remaining IMF-DM degeneracy will require more extensive analysis, we carry out a simple exercise to provide initial clues, inspired by Dutton et al. (2012b). As in that paper, we select the galaxies with mean central stellar surface-densities $\Sigma_* > 2500 M_{\odot} \text{ pc}^{-2}$, and analyze them the same as the full sample. The rationale is that such galaxies are the most star-dominated, and the least sensitive to DM uncertainties. The results are shown by the gray curves in Figure 2, where as expected, the model curves are closer together. The implied δ_{IMF} normalization is fairly low –




Figure 3. Inferred DM fraction within $1 R_e$, vs. σ_e , for the SPIDER sample with different modeling assumptions. The solid and dashed black curves are our default no-AC-NFW and AC-NFW models. The gray curves are for fixed Chabrier IMF, with isothermal and constant- Υ_{dyn} models, bracketing the range of non-parametric models (solid and dotted curves, respectively; see T+12). With standard Λ CDM halos and a variable IMF, f_{DM} is constant or mildly decreasing with σ_e ; with a fixed IMF, f_{DM} is strongly increasing (cf. Thomas et al. 2011 figure 16).

similar to the full-sample AC results. We may then apply this IMF result to the full sample if we assume no additional systematic $\delta_{\text{IMF}}-\Sigma_*$ correlation (cf. Schulz et al. 2010).

Although this paper is primarily concerned with the IMF, we briefly examine some implications for the central DM content. Our main mass models are, by construction, fully consistent with current expectations for Λ CDM halo profiles, while also agreeing with the observations for a plausible IMF range ($\delta_{\text{IMF}} \sim 0.5-2.0$).

We show the implied f_{DM} within $1 R_e$ in Figure 3. We find fairly universal values of $f_{\text{DM}} \sim 0.2$ and ~ 0.5 for the no-AC-NFW and AC-NFW models, respectively. These results are not altered appreciably in the alternative models explored above. Note also that the high- Σ_* test above prefers the AC-NFW model.

The Figure also shows that if we adopted a constant IMF, then we would infer a strong increase of f_{DM} with σ_e . Such behavior has been invoked as the driver for the “tilt” of the ETG fundamental plane (see Tortora et al. 2009 and references therein), but now with renewed comparison to realistic DM halo models, we find that the tilt is driven in part by the IMF. Put differently, the observed $\Upsilon_{\text{dyn}}-\sigma_e$ relation is too steep to explain through standard DM models, and requires an additional factor. Again, the IMF variation is in principle degenerate with systematic variations in AC, which we cannot completely exclude.

4. LITERATURE COMPARISONS

We now inventory other results from the literature, and see if they agree with our SPIDER-based results, showing the comparisons in various panels of Figure 4. We begin with those studies that used similar hybrid approaches,

comparing SPS-mass estimates to total mass using dynamics or lensing. Rather than exhaustively comparing all such results, we will focus on the studies that explicitly derived δ_{IMF} for large samples of low- z ETGs.

The study most closely related to ours is Dutton et al. (2012a), who analyzed SDSS data for ETGs (colors and σ_* , with SPS+Jeans modeling). Their sample was larger but without the Sérsic models and near-infrared photometry from SPIDER, which ought to provide more accurate results. Their results for a no-AC-NFW model are shown as a dotted orange curve in panel (a), which is reassuringly similar to our no-AC-NFW result, with the $\sim 30\%$ residual difference in δ_{IMF} illustrating the level of systematic uncertainties for a fixed dataset and method. The solid line shows their refined result from a multi-parameter fit: they find DM halos that are slightly expanded, and consequently a slightly heavier IMF.

In panel (b) we show results from the SAURON project, using both spectroscopically-based SPS models and detailed two-dimensional Jeans dynamical analyses. For the full ATLAS^{3D} sample, we used the color-coding in figure 2(e) of Cappellari et al. (2012) to regenerate roughly their no-AC-NFW $\delta_{\text{IMF}}-\sigma_e$ results. As a check, we also show their prototype individual galaxy results from Cappellari et al. (2006) (no-DM models), omitting the young galaxies that are thought to suffer from SPS modeling errors. As with SPIDER, the ATLAS^{3D} project found that the overall δ_{IMF} normalization was degenerate to the DM assumptions, but the trend with σ_e was robust. Direct comparison in Figure 4 reveals that the two sets of SAURON results are consistent with SPIDER to within the errors, which is also the case for a similar study of high- z galaxies (Cappellari et al. 2009).

Panel (c) shows results from Thomas et al. (2011) and Wegner et al. (2012), who carried out spectroscopic SPS and detailed orbit-modeling of ETGs found in clusters (including Coma), using a variety of mass models (constant- Υ_{dyn} , no-AC-NFW, and cored halos). There is a systematic offset in δ_{IMF} between the two studies, but overall, the results are consistent with our no-AC-NFW results, both in amplitude and in slope.

Panel (d) shows gravitational lensing results, primarily from the SLACS survey of ETGs (Treu et al. 2010), using no-AC-NFW models as well as color-based SPS models and stellar-dynamics constraints. Their δ_{IMF} normalization agrees well overall with ours, but their slope is somewhat steeper.²

All four hybrid studies from the literature suggest the same basic result as ours: that the IMF of ETGs varies from MW-like at low σ_e to Salpeter-like at high σ_e , modulo some lingering uncertainties from the DM-IMF degeneracy.

In addition to the large-sample studies, we examine a few key single-galaxy results. These include extended kinematics data with NFW-based modeling (Napolitano et al. 2009, 2011), as shown in panel (c), and the “Jackpot” double lens (Sonnenfeld et al. 2012), in panel (d). These results are consistent with the trend from SPIDER.

² Many analyses of the SLACS data have been carried out in the literature, and not all of them are in agreement about the IMF conclusions. It is beyond the scope of this paper to resolve these inconsistencies, but we note that our own analysis in Tortora et al. (2010) agrees with the SLACS-team analysis.




Figure 4. IMF mismatch-parameter vs. stellar velocity dispersion, comparing the SPIDER results (black curves) to various literature studies (see main text and legend, with abbreviations in the reference list). In several panels, typical error-bars are shown to the left. The filled CvD12 datapoint is from four galaxies’ stacked spectra. There is generally good consistency in the inferred IMF trends from the different studies.

Our culminating comparisons are with a completely different set of results, based purely on modeling of IMF-sensitive spectral lines (see Section 1). Most of these lines are susceptible to degeneracies with elemental abundances (e.g., sodium or calcium), and we consider only those studies that have directly accounted for such effects.

We first show in panel (e) the results from CvD12, who studied a subset of the SAURON sample. They fitted spectral features across a wide wavelength range, focusing on the IMF indicators Na I, Ca II, and the Wing-Ford band, while adopting a broken-power-law IMF form and fitting for the relevant elemental abundances. Their inferred δ_{IMF} values turn out to agree well with both the normalization and the trend versus σ_e from the SPIDER results (for the no-AC-NFW models in particular)³. Recalling that the high δ_{IMF} values from hybrid studies could be due to either a bottom-heavy (extra dwarfs) or a top-heavy (extra remnants) IMF, the CvD12 results

³ CvD12 compared their Υ_* results to *total* dynamical values from SAURON in order to check that they did not violate those constraints. However, they did not compare to *decomposed* dynamical Υ_* inferences for consistency as we do here.

agree with only the first solution.

Note that the apertures probed here are different: $R_e/8$ for CvD12, and $0.3-0.7 R_e$ for SPIDER (decreasing with σ_*). The close agreement of the results on average thus implies that the IMF does not vary spatially on these scales, or that the AC model is the correct solution, and δ_{IMF} decreases with galactocentric radius (which is plausible).

Panel (f) shows results from Smith et al. (2012), who studied a large sample of Coma-cluster ETGs. They used the same SPS models as CvD12 to fit near-infrared spectroscopic line-indices, analyzing the Wing-Ford band and the Ca I line separately. The former line was susceptible to uncertainties in the Na abundance, but not the latter. We have converted their results to inferred δ_{IMF} by straightforward interpolation between the Chabrier, Salpeter, and $\alpha = 3$ models in their figure 10. The final results are somewhat noisy and uncertain, but imply an overall Salpeter-like normalization, and no obvious trend with σ_e . This agrees with all the aforementioned results at high- σ_e , but not at low- σ_e . It is possible that these Coma-cluster galaxies are genuinely different. We see very little dependence of δ_{IMF} on environment in SPI-

DER, but we also have very few cluster galaxies in our sample.

Spiniello et al. (2012) analyzed NaD, Na I, and TiO₂ lines in SDSS spectra of ETGs with $\sigma_\star > 200 \text{ km s}^{-1}$, comparing to the same SPS models as CvD12. They inferred a Salpeter-like IMF at low $\sigma_\star$, which is consistent with SPIDER. At high $\sigma_\star$, they inferred $\alpha \sim 3$, which would imply $\delta_{\text{IMF}} \sim 4$ and violate the total mass constraints both from SPIDER and from the lens galaxy that these authors also studied. This conflict suggests further work is needed on the line-index modeling.

An ideal comparison with our SPIDER results would be the work of Ferreras et al. (2012), who analyzed Na I line-strengths from the SPIDER parent dataset. Qualitatively, their results imply a similar increase of δ_{IMF} with σ_e , but more quantitative inferences will have to await more detailed analyses that allow for Na enhancement.

It would also be fascinating to see whether or not spiral bulges follow the same $\delta_{\text{IMF}}-\sigma_\star$ trends as ETGs. Falcón-Barroso et al. (2003) found that bulges showed anti-correlations between $\sigma_\star$ and Ca II line-strengths, similar to ETGs, but they could not determine if this was an IMF effect. We have attempted to investigate further, using δ_{IMF} inferences for bulges from the literature (e.g., de Blok et al. 2008; CvD12; Dutton et al. 2012c). We find hints of an ETG-like $\delta_{\text{IMF}}-\sigma_\star$ correlation for the bulges, but we also find the same for the disks, which would be peculiar. The apparent scatter for both bulges and disks is enormous, suggesting more work is needed to address the systematic errors, and to understand any additional trends with detailed morphology.

5. DISCUSSION AND CONCLUSIONS

We have analyzed the dynamics and stellar populations of a large sample of ETGs from the SPIDER project, and found compelling evidence for heavier IMFs in the central regions of higher- $\sigma_\star$ galaxies. The IMF mismatch relative to Chabrier is $\delta_{\text{IMF}} \sim 0.5-1.1$ at $\sigma_\star \sim 125 \text{ km s}^{-1}$, and $\sim 1.2-1.8$ at $\sim 250 \text{ km s}^{-1}$. The $\delta_{\text{IMF}}-\sigma_\star$ trend is robust to different modeling assumptions, and accounts for much of the tilt in the fundamental plane. The distribution of DM is degenerate with the overall IMF normalization and difficult to constrain.

We have performed the first general inventory of IMF results from a variety of studies in the literature, using both pure and hybrid techniques, and found that these generally agree well with our SPIDER results. There is remarkably widespread agreement on a Salpeter-like IMF for massive ETGs ($\sigma_e \gtrsim 200 \text{ km s}^{-1}$). At lower σ_e , the data are still somewhat limited and the different results have not yet converged, but most of the studies point to MW-like IMFs. Agreement on a super-Salpeter IMF in the most massive galaxies ($\sigma_e \gtrsim 275 \text{ km s}^{-1}$) is also not yet universal.

These results appear to be fully compatible with underlying Λ CDM halos. However, more detailed conclusions about halo contraction or expansion are still elusive, and the data on galaxy centers do not clearly rule out alternative DM models once the variable IMF is accounted for.

More work is clearly needed to understand the systematics in the different analyses; to build up better statistics on a wide range of galaxy types, environments, and red-

shifts; and to determine which parameters correlate best with IMF variations (e.g., metallicity or starburst intensity; CvD12; Smith et al. 2012). It may also be particularly helpful to venture beyond the centers of galaxies, using data from a wide baseline in radius to help break the IMF-DM degeneracies.

It appears we are nearing convergence on determining *what* the basic components of galaxies are (distributions of stars and DM). The next challenge will be to understand *why* these arrive at their distributions. What drives the power-spectrum in cloud fragmentation and star formation? How do baryonic processes interact with and re-shape their surrounding DM halos?

We thank Matt Auger, Michele Cappellari, Charlie Conroy, Reinaldo de Carvalho, Aaron Dutton, Francesco La Barbera, and Tommaso Treu for helpful discussions. CT was supported by the Swiss National Science Foundation and the Forschungskredit at the University of Zurich. AJR was supported by the National Science Foundation Grant AST-0909237.

REFERENCES

- Abadi, M. G., Navarro, J. F., Fardal, M., Babul, A., & Steinmetz, M. 2010, *MNRAS*, 407, 435 (A+10)
- Auger, M. W., Treu, T., Gavazzi, R., et al. 2010, *ApJ*, 721, L163
- Bastian, N., Covey, K. R., & Meyer, M. R. 2010, *ARA&A*, 48, 339
- Bell, E. F., & de Jong, R. S. 2001, *ApJ*, 550, 212
- Bershady, M. A., Martinsson, T. P. K., Verheijen, M. A. W., et al. 2011, *ApJ*, 739, L47
- Blumenthal, G. R., Faber, S. M., Flores, R., & Primack, J. R. 1986, *ApJ*, 301, 27 (B+86)
- Brewer, B. J., Dutton, A. A., Treu, T., et al. 2012, *MNRAS*, 422, 3574
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Burkert, A. 1995, *ApJ*, 447, L25
- Cappellari, M., Bacon, R., Bureau, M., et al. 2006, *MNRAS*, 366, 1126 (C+06)
- Cappellari, M., di Serego Alighieri, S., Cimatti, A., et al. 2009, *ApJ*, 704, L34
- Cappellari, M., McDermid, R. M., Alatalo, K., et al. 2012, *Nature*, 484, 485 (C+12)
- Cenarro, A. J., Gorgas, J., Vazdekis, A., Cardiel, N., & Peletier, R. F. 2003, *MNRAS*, 339, L12
- Chabrier, G. 2003, *PASP*, 115, 763
- Conroy, C., & van Dokkum, P. 2012, *ApJ*, submitted, arXiv:1205.6473 (CvD12)
- de Blok, W. J. G., Walter, F., Brinks, E., et al. 2008, *AJ*, 136, 2648
- Dutton, A. A., Maccio', A. V., Mendel, J. T., & Simard, L. 2012a, *MNRAS*, submitted, arXiv:1204.2825
- Dutton, A. A., Mendel, J. T., & Simard, L. 2012b, *MNRAS*, 422, L33
- Dutton, A. A., Treu, T., Brewer, B. J., et al. 2012c, *MNRAS*, submitted, arXiv:1206.4310
- Falcón-Barroso, J., Peletier, R. F., Vazdekis, A., & Balcells, M. 2003, *ApJ*, 588, L17
- Ferreras, I., La Barbera, F., de Carvalho, R. R., et al. 2012, *ApJ*, submitted, arXiv:1204.3823
- Ferreras, I., Saha, P., & Burles, S. 2008, *MNRAS*, 383, 857
- Gnedin, O. Y., Kravtsov, A. V., Klypin, A. A., & Nagai, D. 2004, *ApJ*, 616, 16 (G+04)
- Grillo, C. 2010, *ApJ*, 722, 779
- Grillo, C., & Gobat, R. 2010, *MNRAS*, 402, L67
- Kroupa, P. 2001, *MNRAS*, 322, 231
- La Barbera, F., de Carvalho, R. R., de La Rosa, I. G., et al. 2010, *MNRAS*, 408, 1313
- Macciò, A. V., Dutton, A. A., & van den Bosch, F. C. 2008, *MNRAS*, 391, 1940
- Milgrom, M. 1983, *ApJ*, 270, 371

- Moster, B. P., Somerville, R. S., Maulbetsch, C., et al. 2010, *ApJ*, 710, 903 (M+10)
- Napolitano, N. R., Romanowsky, A. J., & Tortora, C. 2010, *MNRAS*, 405, 2351
- Napolitano, N. R., Romanowsky, A. J., Coccato, L., et al. 2009, *MNRAS*, 393, 329 (N+09)
- Napolitano, N. R., Romanowsky, A. J., Capaccioli, M., et al. 2011, *MNRAS*, 411, 2035 (N+11)
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, *ApJ*, 462, 563
- Renzini, A., & Ciotti, L. 1993, *ApJ*, 416, L49
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Schneider, A., Smith, R. E., Macciò, A. V., & Moore, B. 2012, *MNRAS*, 424, 684 (S+12a)
- Schulz, A. E., Mandelbaum, R., & Padmanabhan, N. 2010, *MNRAS*, 408, 1463
- Smith, R. J., Lucey, J. R., & Carter, D. 2012, *MNRAS*, submitted, arXiv:1206.4311 (S+12b)
- Sonnenfeld, A., Treu, T., Gavazzi, R., et al. 2012, *ApJ*, 752, 163 (S+12c)
- Spiniello, C., Koopmans, L. V. E., Trager, S. C., Czoske, O., & Treu, T. 2011, *MNRAS*, 417, 3000
- Spiniello, C., Trager, S. C., Koopmans, L. V. E., & Chen, Y. P. 2012, *ApJ*, 753, L32
- Suyu, S. H., Hensel, S. W., McKean, J. P., et al. 2012, *ApJ*, 750, 10
- Thomas, J., Saglia, R. P., Bender, R., et al. 2011, *MNRAS*, 415, 545 (T+11)
- Tortora, C., La Barbera, F., Napolitano, N. R., de Carvalho, R. R., & Romanowsky, A. J. 2012, *MNRAS*, in press, arXiv:1201.2945 (T+12)
- Tortora, C., Napolitano, N. R., Romanowsky, A. J., Capaccioli, M., & Covone, G. 2009, *MNRAS*, 396, 1132
- Tortora, C., Napolitano, N. R., Romanowsky, A. J., & Jetzer, P. 2010, *ApJ*, 721, L1
- Treu, T., Auger, M. W., Koopmans, L. V. E., et al. 2010, *ApJ*, 709, 1195 (T+10)
- van Dokkum, P. G., & Conroy, C. 2010, *Nature*, 468, 940
- Wegner, G. A., Corsini, E. M., Thomas, J., et al. 2012, *AJ*, in press, arXiv:1206.5768 (W+12)